

B-tagging plots for Moriond

Dataset

- All p10.14.00 data between runs 142028 and 143769
- E/F Muon runs removed, bad calorimeter runs removed
- Selected on tight muon + loose jet and *mulptxxctxx_CJT3* trigger
- 8337 events, 183 nb^{-1}

Muon + jet cross section




Efficiencies


- Muon Acceptance: 0.89
- Muon L1 trigger efficiency: 0.4
- Muon offline reconstruction & ID: 0.33


- Jet L1 trigger efficiency: 0.98
- Jet offline reconstruction & ID: 0.98

Jet Energy Scale applied to energy of jets


Muon momentum corrected for energy loss in calorimeter







DØ Run 2 Preliminary



B-jet content measurement using $P_{T\text{Rel}}$

- Same data sample
- No p_T cut on muon, no p_T cut on jet
- No η cuts
- $\delta R(\text{jet}, \mu) < 0.7$

Muon direction updated with scintillator hit

MC samples:

bbbar: SAM project bbbar_p100801, 117500 events

ccbar: SAM project ccbar_p100801, 50000 events


QCD: SAM projects lancs_qcd_131, nikhef_qcd_131, 1.1M events

Data muon momentum resolution worse than MC momentum resolution → MC muon momentum smeared with factor 0.09

$P_{T\text{Rel}}$ is measured with respect to new combined muon + jet axis

Plots for both local muons and global muons available

DØ Run 2 Preliminary



DØ Run 2 Preliminary

